

The asymptotic number of rooted trees in a tree*

Xueliang Li, Yiyang Li, Yongtang Shi

Center for Combinatorics and LPMC-TJKLC

Nankai University, Tianjin 300071, China

Email: lxl@nankai.edu.cn, liycldk@mail.nankai.edu.cn, shi@nankai.edu.cn

Abstract

Let $\mathcal{T}_n$ be the set of trees with n vertices. Suppose that each tree in $\mathcal{T}_n$ is equally likely. We show that the number of rooted trees corresponding to any given tree equals $(\mu_r + o(1))n$ in almost every tree of $\mathcal{T}_n$, where μ_r is a constant. As an application, we show that the number of any given pattern in $\mathcal{T}_n$ is also asymptotically normally distributed with mean $\sim \mu_M n$ and variance $\sim \sigma_M n$, where μ_M, σ_M are some constants related to the given pattern.

Keywords: tree; rooted tree; pattern; generating function; limiting distribution; automorphism

AMS Subject Classification (2010): 05C05, 05C30, 05D40, 05A15, 05A16.

1 Introduction

A *pattern* $\mathcal{M}$ is a given small tree. We say that $\mathcal{M}$ *occurs* in a tree T if $\mathcal{M}$ is a subtree of T in the sense that the degree of each internal vertex (of degree more than one) of $\mathcal{M}$ matches the degree of the corresponding vertex in T , while each external vertex (of degree one) of $\mathcal{M}$ matches a vertex of T with an arbitrary degree. Occasionally, we say that the pattern is in a tree instead of that the pattern occurs in a tree for abbreviate. Let $\mathcal{T}_n$ be the set of trees with n vertices. If we use $X_{n,M}$ to denote the number of a given pattern $\mathcal{M}$ in $\mathcal{T}_n$, then $X_{n,M}$ is a random variable with probability

$$Pr(X_{n,M} = k) = \frac{t_{n,k}}{t_n},$$

where $t_{n,k}$ denotes the number of such trees in $\mathcal{T}_n$ that the number of pattern $\mathcal{M}$ in each of the trees is k , and $t_n = |\mathcal{T}_n|$.

*Supported by NSFC.

Moreover, let $\mathcal{R}_n$ be the set of rooted trees. We can also consider the number of a given pattern in $\mathcal{R}_n$. We still use $X_{n,M}$ to denote the random variable in $\mathcal{R}_n$, which would not make any ambiguity.

The main work of this paper is to show that some random variable satisfies

$$\frac{Y_n - E(Y_n)}{\sqrt{\text{Var}(Y_n)}} \rightarrow_w \mathcal{N}(0, 1),$$

where $\mathcal{N}(0, 1)$ is the random variable with standard normal distribution and $\rightarrow_w$ means weak convergence. We then call this Y_n *asymptotically normal*. Moreover, if

$$\frac{Y_n - \mu n}{\sqrt{\sigma n}} \rightarrow_w \mathcal{N}(0, 1),$$

then Y_n is asymptotically normal with mean $\sim \mu n$ and variance $\sim \sigma n$. We refer the readers to [7] for details.

In fact, it was shown in [2] that in $\mathcal{R}_n$ the number $X_{n,M}$ of any given pattern is asymptotically normal with mean $\sim \mu_M n$ and variance $\sim \sigma_M n$, where μ_M and σ_M are some constants corresponding to the given pattern. But, for the set $\mathcal{T}_n$ there is no such a result on normal distribution. In [6], the authors proved that for any given pattern in $\mathcal{T}_n$ the limiting distribution has a density $(A + Bt^2)e^{Ct^2}$, where A, B, C are some constants. The mean and variance of the number of any given pattern are still asymptotically $\mu_M n$ and $\sigma_M n$ where the constants are the same as in $\mathcal{R}_n$. Clearly, if one shows that $B = 0$, then the distribution is normal. For some special patterns, such as a star (or a node with a given degree) pattern [4], a double-star pattern [8], and a path pattern [7], the corresponding limiting distributions were proved to be normal. For some previous work we refer to Robinson and Schwenk [11]. For more details, we refer to [2, 6, 7, 11]. However, for any given pattern Kok in his thesis [7] claimed that it seems much more difficult to demonstrate the normality. In this paper, we will solve this problem from a new point of view which is different from the existing ones. We study the number of rooted trees in a tree and get that for almost every tree the number of corresponding rooted trees is $(\mu_r + o(1))n$. Then, for any given pattern $\mathcal{M}$, since we already knew that the number of pattern $\mathcal{M}$ in $\mathcal{R}_n$ is asymptotically normal, as a consequence, in $\mathcal{T}_n$ the limiting distribution for $\mathcal{M}$ is also normal.

We organize this paper as follows. In Section 2, we will introduce some basic knowledge that will be used in our proofs. In Section 3, we will present the detailed proofs. We concentrate on the number of rooted trees in a tree. Section 4 is devoted to study the limiting distribution for any given pattern.

2 Preliminaries

Analogous to patterns, for each tree we use X_n to denote the number of corresponding rooted trees. Clearly, X_n is also a random variable in $\mathcal{T}_n$. We introduce the following two functions:

$$t(x) = \sum_{n \geq 1} t_n x^n,$$

$$t(x, u) = \sum_{n \geq 1, k \geq 0} t_{n,k} x^n u^k,$$

where the coefficient $t_{n,k}$ denotes the number of such trees in $\mathcal{T}_n$ that each of the trees has k rooted trees. Clearly, $\sum_{k \geq 0} t_{n,k} = t_n$. We always assume that every tree of $\mathcal{T}_n$ is equally likely. Then, $\Pr(X_n = k) = \frac{t_{n,k}}{t_n}$.

Let T_n be a tree in $\mathcal{T}_n$. An automorphism Φ of T_n is defined as

$$\begin{aligned} \Phi : v_i &\rightarrow \Phi(v_i) \\ v_i v_j &\rightarrow \Phi(v_i) \Phi(v_j) \end{aligned}$$

where v_i and v_j are any two vertices in T_n and $v_i v_j$ is an edge of T_n joining vertices v_i and v_j . We call that two vertices u and v of T_n are in the same vertex class if u can be mapped to v by some automorphism. Clearly, this sets up an equivalent relation on the vertex set of T_n , and hence the vertices in T_n are partitioned into some classes. If we designate every vertex in a same vertex class to be the root, we shall get the same rooted tree. Exactly to say, suppose that v_i is mapped to v_j under an automorphism Φ of T_n . One can illustrate that the automorphism Φ is also an isomorphism that maps the tree rooted at v_i to the tree rooted at v_j just by definition.

Hence, the number of rooted trees in a tree is exactly the number of vertex classes in the tree. Then, let X_n also represents the number of vertex classes in the tree under automorphisms. Therefore, we can introduce X_n on the rooted tree space $\mathcal{R}_n$.

If we consider X_n in $\mathcal{R}_n$, we also suppose that each tree in $\mathcal{R}_n$ is equally likely. We can define similar generating functions on $\mathcal{R}_n$, and let $r(x)$, $r(x, u)$ be the related functions, respectively. One can see that $r(x, 1) = r(x)$. Suppose

$$r(x, u) = \sum_{n \geq 1, k \geq 0} r_{n,k} x^n u^k,$$

where $r_{n,k}$ is the number of rooted trees in $\mathcal{R}_n$ that have k vertex classes. It follows that in $\mathcal{R}_n$,

$$\Pr(X_n = k) = \frac{r_{n,k}}{r_n},$$

where $r_n = |\mathcal{R}_n|$.

We should notice that when we count the number of vertex classes in a rooted tree, the root itself always forms a class with a single vertex, since any automorphism on a rooted tree must map the root to itself. That is a bit different from the case for trees.

Furthermore, suppose that the radius of convergence of $r(x)$ is x_0 . Otter [9] showed that x_0 satisfies that $r(x_0) = 1$ and the asymptotic expansion of $r(x)$ is

$$r(x) = 1 - b_1(x_0 - x)^{1/2} + b_2(x_0 - x) + b_3(x_0 - x)^{3/2} + \dots, \quad (1)$$

where $x_0 \approx 0.3383219$ and $b_1 \approx 2.6811266$. And, $t(x)$ has a similar expansion, namely,

$$t(x) = c_0 + c_1(x_0 - x) + c_2(x_0 - x)^{3/2} + \dots. \quad (2)$$

Applying approximate theory [10] on Eqs.(1) and (2), we get that

$$\begin{aligned} t_n &\sim \frac{Cx_0^{-n}}{n^{5/2}}, \\ r_n &\sim \frac{Dx_0^{-n}}{n^{3/2}}, \end{aligned}$$

where C and D are some constants. We refer the readers to [11]. It has been showed that $C = 0.5349\dots$ and $D = 0.4399\dots$. The approximate theory used in [11] is due to Pólya. The original paper [10] gives us more details.

In what follows, we first investigate X_n in $\mathcal{R}_n$. To start with, we need the following two lemmas. We refer the readers to [3, 7] for detailed information.

Lemma 1. *Suppose that $F(x, y, u)$ is an analytic function around at $(x_0, y_0, 1)$ such that $F(x_0, y_0, 1) = y_0$, $F_{y_0}(x_0, y_0, 1) = 1$, $F_{yy}(x_0, y_0, 1) \neq 0$ and $F_x(x_0, y_0, 1) \neq 0$. Then there exist a neighborhood U_0 of $(x_0, 1)$, a neighborhood U_1 of y_0 and analytic functions $g(x, u)$, $h(x, u)$ and $f(u)$ which are defined on U_0 such that the only solutions $y \in U_1$ with $y = F(x, y, u)$ and $(x, u) \in U_0$ are given by $y(x, u) = g(x, u) + h(x, u)\sqrt{1 - \frac{x}{f(u)}}$. Furthermore, $g(x_0, 1) = y_0$ and $h(x_0, 1) = \sqrt{\frac{2f(1)F_x(x_0, y_0, 1)}{F_{yy}(x_0, y_0, 1)}}$. $\square$*

Lemma 2. *Let $y(x, u)$ denote the function defined on a neighborhood U of $(x_0, 1)$, and $y(x, u) = F(x, y(x, u), u) = g(x, u) + h(x, u)\sqrt{1 - \frac{x}{f(u)}}$. If $y(x, 1)$ is aperiodic, i.e., if from $y(x, 1) = x^r \tilde{y}(x^d, 1)$ with some power series $\tilde{y}(x, u)$ follows that $d = 1$ and all the Taylor coefficients of $F_y(x, y, u)$ are non-negative, then there exists an $\eta > 0$ such that $y(x, u)$ can be analytically continued in*

$$\tilde{U} = \{(x, u) : |x| < x_0 + \eta, |u| < 1 + \eta, \arg(x - f(u)) \neq 0, x \neq f(u)\}.$$

Moreover, if $y(x, u) = \sum y_{n,k} x^n u^k = \sum [y_n(u)] x^n$ and $y_{n,k} \geq 0$, then

$$y_n(u) = \frac{h(f(u), u)}{2\sqrt{\pi}n^{3/2}} f(u)^{-n} + O\left(\frac{f(u)^{-n-1}}{n^{5/2}}\right).$$

And if $h(f(1), 1) \neq 0$, we get that X_n is asymptotically normal with mean $\sim \mu n$ and variance $\sim \sigma n$. $\square$

Remark: In [3] and [7], the authors always assumed that all the Taylor coefficients of $F(x, y, u)$ are non-negative. But, we can extend this condition in the proofs in [3] and [7] to the following one: all the Taylor coefficients of $F_y(x, y, u)$ are non-negative.

3 The number of rooted trees in a tree

Now we concentrate on the number of vertex classes of a rooted tree. Recalled that an automorphism of a rooted tree must map the root to itself, which is a bit different from an automorphism of a tree, namely, the root always forms a vertex class with a single vertex. We shall show that X_n is asymptotically normal with mean $(\mu_r + o(1))n$ and variance $(\sigma_r + o(1))n$ in $\mathcal{R}_n$.

In what follows, there appears an expression of the form $Z^*(S_n; f(x, u))$ (or $Z(S_n; f(x))$), which is the substitution of the counting series $f(x, u)$ (or $f(x)$) into the cycle index $Z(S_n)$ of the symmetric group S_n . This involves replacing each variable s_i in $Z(S_n)$ by $f(x^i, u)$ (or $f(x^i)$). For instance, if $n = 3$, then $Z(S_3) = (1/3!)(s_1^3 + 3s_1s_2 + 2s_3)$ and $Z(S_3; f(x)) = (1/3!)(f(x)^3 + 3f(x)f(x^2) + 2f(x^3))$, $Z^*(S_3; f(x, u)) = (1/3!)(f(x, u)^3 + 3f(x, u)f(x^2, u^2) + 2f(x^3, u^3))$. We refer the readers to [5] for details. And in [5], it was shown that

$$r(x) = x \cdot \sum_{n \geq 0} Z(S_n; r(x)) = x \cdot e^{\sum_{k=1} \frac{r(x^k)}{k}}.$$

The coefficient of x^p in $Z(S_n; r(x))$ is the number of rooted trees of order $p + 1$ whose roots have degree n . Multiplication of $Z(S_n; r(x))$ by x corrects the power of x so that x^p in $xZ(S_n; r(x))$ is the number of those trees with p vertices. This expression $Z(S_n; r(x))$ follows from Pólya Enumeration Theorem [5].

Analogously, we take the same procedure for $r(x, u)$ in this paper. But, here we should notice that if the same two copies of a rooted tree with k vertex classes connect to a root, then the number of vertex classes of the new rooted tree is $k + 1$, because there is only one new class, i.e., the new root, which is different from the procedure for calculating the number of a given pattern. Hence, we use $r(x^k, u)$ to denote the generating function for k copies of a rooted tree. And we can get

$$r(x, u) = xu \cdot e^{\sum_{k \geq 1} \frac{1}{k} r(x^k, u)} + \psi(x, u), \quad (3)$$

where the modification term $\psi(x, u)$ is a polynomial.

For instance, suppose that the tree has a root of degree 2. Then, $xu \cdot Z^*(S_2, r(x, u)) = xu \cdot \frac{1}{2}(r(x, u)^2 + r(x^2, u))$. We have $r_{n,k}$ choices to form a rooted tree with the same two

branches. In $r(x, u)^2$ the term $r_{n,k}x^{2n}u^{2k}$ denotes the number of such rooted trees. We should notice that the power $2k$ must be corrected into k which means that the number of vertex classes is still k . But, in $r(x^2, u)$ the term $r_{n,k}x^{2n}u^k$ denotes the number of those rooted trees with two same branches. Moreover, note that $r(x^2, u)$ and $r(x^2, u^2)$ are polynomials, since $r(x, u)$ is a polynomial. Hence, there must be a polynomial modification term $\psi_2(x, u)$ such that $xu \cdot Z^*(S_2, r(x, u)) + \psi_2(x, u)$ performs the generating function of the trees with roots of degree 2. We can see that for any number of branches, we have to modify the function when counting the numbers corresponding to the cases that some branches are the same. Hence, in general, expression (3) follows.

Let $y = r(x, u)$, and $F(x, y, u) = xu \cdot e^{\sum_{k \geq 1} \frac{1}{k} r(x^k, u)} + \psi(x, u)$. Here, the Taylor coefficients of x, y and u may not be non-negative. Recall that $r(x, 1) = r(x)$, that is, $\psi(x, 1) = 0$. Recall also that there exists a real number x_0 such that $r(x_0) = 1$ and x_0 is the convergence radius.

Then, we have $F(x_0, y(x_0, 1), 1) = 1 = y(x_0, 1) = r(x_0, 1)$ and

$$F_y(x, y, u) = xu \cdot e^{\sum_{k \geq 1} \frac{1}{k} r(x^k, u)},$$

which implies that $F_y(x_0, y_0, 1) = 1$. Moreover, $F_{yy}(x_0, y_0, 1) \neq 0$ and $F_x(x_0, y_0, 1) \neq 0$. That is, all the conditions of Lemma 1 hold. Furthermore, we have that the Taylor coefficients of $F_y(x, y, u)$ are non-negative, because in F_y all $r(x^k, u)$'s ($k \geq 2$) are polynomials and the expression of F_y has an exponential form. Thus, by Lemma 2 we have that the random variable X_n is asymptotically normal with mean

$$E(X_n) \sim \mu_r n \quad (n \rightarrow \infty)$$

and variance

$$\text{Var}(X_n) \sim \sigma_r n \quad (n \rightarrow \infty),$$

where μ_r and σ_r are some constants.

In this paper, we mainly focus on the overall property of a probability space. Following the book [1], we will say that *almost every* (a.e.) graph in a graph space $\mathcal{G}_n$ has a certain property Q if the probability $\Pr(Q)$ in $\mathcal{G}_n$ converges to 1 as n tends to infinity. Occasionally, we will say *almost all* instead of almost every.

From Chebyshev inequality

$$\Pr[|X_n - E(X_n)| > n^{3/4}] \leq \frac{\text{Var} X_n}{n^{3/2}} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

it follows that for almost all rooted trees, $E(X_n) + n^{3/4} \leq X_n \leq E(X_n) + n^{3/4}$, namely, $X_n = (1 + o(1))E(X_n)$. We can get that

Theorem 3. *For almost all rooted trees in $\mathcal{R}_n$, the number of vertex classes under automorphisms is $(\mu_r + o(1))n$. $\square$*

Therefore, we can study the number of vertex classes in a tree. To get the final result, we need another property as follows. We have defined the number of vertex classes of a tree. And, we call a vertex *fixed* if this single vertex forms a class.

Lemma 4. *Almost every tree in $\mathcal{T}_n$ has more than $\frac{1}{24}n$ fixed vertices.*

Proof. We prove this result by contradiction. Suppose that $\mathcal{T}'_n$ is a subset of $\mathcal{T}_n$ such that every tree T'_n in $\mathcal{T}'_n$ has at most $\frac{1}{24}n$ fixed vertices. We first show that these fixed vertices form a subtree in T'_n . In fact, for any two fixed vertices v_1 and v_2 , they can only map to v_1 and v_2 among themselves, respectively. Thus, any (v_1, v_2) -path maps to the (v_1, v_2) -path under any automorphism. So, all vertices in a (v_1, v_2) -path are fixed ones, that is, all the fixed vertices form a connected subgraph of T'_n . Consequently, the fixed vertices induce a subtree T''_n of T'_n and $|T''_n| \leq \frac{1}{24}n$. If $|T''_n| = 0$, then the tree T'_n has a symmetrical edge. The structure of T'_n is determined by one half of the vertices in T'_n . Hence, the number of trees in $\mathcal{T}_n$ having a symmetrical edge is at most $|\mathcal{T}_{\frac{n}{2}}|$, and so $\frac{|\mathcal{T}_{\frac{n}{2}}|}{|\mathcal{T}_n|} \rightarrow 0$.

Then, we always suppose $|T''_n| > 0$. Let u be a vertex in T''_n . Suppose that H_u is a subtree of T'_n attaching to u such that all the vertices in H_u are not in T''_n . Suppose there are m copies of H_u after deleting u . We have $m \geq 2$, otherwise the vertex in H_u connecting to u is also a fixed vertex, which is a contradiction. If m is even, we get rid off $m/2$ copies of H_u . If m is odd, we delete $(m+1)/2$ copies of H_u . We repeat this operation on all vertices in T''_n . At the end, this produces a new tree A with at most $\frac{1}{2}(n + 1/24 \cdot n)$ vertices, and we denote the set of these new trees by $\mathcal{A}_{\frac{25}{48}n}$. Moreover, if we replace these $\lfloor \frac{m}{2} \rfloor$ copies of H_u by a vertex, that is, we add some vertices to u and different kinds of H_u correspond to different vertices. Thus, we construct another tree A' . Observe that T''_n is a subtree of A' . We shall show that A' has at most $n/3$ vertices. Color the vertices in A' corresponding to T''_n by red and the others by green. We already knew that there are at most $\frac{1}{24}n$ red vertices. Let u be a red vertex. If a green vertex connecting to u in A' represents just only one vertex of A , then there is only one such vertex connecting to u . One can see an example in Figure 1. Hence, in A' there are at most $\frac{1}{24}n$ vertices representing the subtrees having a single vertex and at most $\frac{1}{2} \cdot \frac{1}{2}n$ vertices representing the other subtrees (or forests) having at least two vertices. Consequently, we get that A' has at most $n/3$ vertices. Moreover, we need the fact that the order of $\mathcal{T}_n$ is asymptotically $\frac{C \cdot x_0^{-n}}{n^{5/2}}$. So, the number of trees with at most n vertices is asymptotically less than $2 \cdot Cx_0^{-n}$.

In the above, we have built a map from $\mathcal{T}'_n$ to $\mathcal{A}_{\frac{25}{48}n}$. Suppose $A_{\frac{25}{48}n}$ is a tree in $\mathcal{A}_{\frac{25}{48}n}$. Then $|A_{\frac{25}{48}n}|$ is at most $\frac{25}{48}n$. The ways to choose k vertices to form a subtree of $A_{\frac{25}{48}n}$ is at most $\binom{\frac{25}{48}n}{k}$. We color these vertices in $A_{\frac{25}{48}n}$ by red. Notice that any tree in $\mathcal{T}''_n$




Figure 1: An example of A' and A .

has at most $\frac{1}{24}n$ vertices. Then, the number of subtrees in $A_{\frac{25}{48}n}$ having at most $\frac{1}{24}n$ vertices is less than $\binom{\frac{25}{48}n}{k} \leq \binom{\frac{25}{48}n}{\frac{1}{24}n}$.

Let T_n'' be one of the subtrees with k vertices. The red vertices have been selected, we suppose that A' is the corresponding tree defined as above. For $u \in T_n''$, each green vertex in A' connecting to u corresponds to a kind of subtree H_u and the number of H_u can be odd or even. That is, for each H_u , we have at most two choices to form a tree with n vertices. Since the number of green vertices is less than $|V(A')| \leq n/3$, there are less than $n/3$ vertices in A' but T_n'' . We can get that there exist at most $2^{\frac{n}{3}}$ trees in $\mathcal{T}_n'$ mapping to $A_{\frac{25}{48}n}$.

Therefore, for trees in $\mathcal{T}_n''$ with k vertices, at most $2^{\frac{n}{3}} \cdot 2C \cdot x_0^{-\frac{25}{48}n} \cdot \binom{\frac{25}{48}n}{k}$ trees in $\mathcal{T}_n'$ map to them. Recall that each T_n' corresponds to some T_n'' . Then we have

$$\begin{aligned} |\mathcal{T}_n'| &\leq \sum_{k=1}^{1/24n} 2^{\frac{n}{3}} \cdot 2C \cdot x_0^{-\frac{25}{48}n} \cdot \binom{\frac{25}{48}n}{k} \\ &\leq \frac{1}{24} n 2^{\frac{n}{3}} \cdot 2C \cdot x_0^{-\frac{25}{48}n} \cdot \binom{\frac{25}{48}n}{\frac{1}{24}n} \\ &= \frac{C}{12} n \cdot 2^{\frac{n}{3}} \cdot x_0^{-\frac{25}{48}n} \cdot \binom{\frac{25}{48}n}{\frac{1}{24}n}. \end{aligned}$$

By Stirling's approximation, i.e., $\frac{n!}{\sqrt{2\pi n}(\frac{n}{e})^n} \rightarrow 1$ as $n \rightarrow \infty$, we can get that when n is large enough,

$$\binom{\frac{25}{48}n}{\frac{1}{24}n} < \frac{C_0}{\sqrt{n}} 1.2^n,$$

where C_0 is a constant. Then,

$$|\mathcal{T}_n'| < C_1 n^{1/2} \cdot 2^{\frac{n}{3}} x_0^{-\frac{25}{48}n} 1.2^n,$$

where C_1 is some real number for large n . It is known that $|\mathcal{T}_n| \sim \frac{C \cdot x_0^{-n}}{n^{\frac{5}{2}}}$. Recall that $x_0 \approx 0.3383219$. Consequently, $\frac{|\mathcal{T}_n'|}{|\mathcal{T}_n|} \rightarrow 0$.

Hence, in conclusion, we get that almost all trees do not belong to $\mathcal{T}'_n$. Then, the result is relevant. $\square$

Next, we proceed to estimate the number of rooted trees for a given tree from Theorem 3 and Lemma 4. The following theorem is established.

Theorem 5. *For almost all trees in $\mathcal{T}_n$, the number of corresponding rooted trees is $(\mu_r + o(1))n$.*

Proof. By Lemma 4, we know that almost every tree has at least $\frac{1}{24}n$ fixed vertices, and denote these trees by $\mathcal{T}_n^*$. Clearly, $\mathcal{T}_n^* \subseteq \mathcal{T}_n$ and $\frac{|\mathcal{T}_n^*|}{|\mathcal{T}_n|} \rightarrow 1$. Let T be a tree in $\mathcal{T}_n^*$. If we pick out one of the fixed vertices to be the root, we can get a rooted tree having the same number of vertex classes. There are at least $\frac{1}{24}n$ rooted trees in which the roots of the rooted trees correspond to the fixed vertices of T . And the number of vertex classes equals to that in T . Hence, there are at least $|\mathcal{T}_n^*| \cdot \frac{1}{24}n$ rooted trees in $\mathcal{R}_n$ such that the roots are fixed vertices in the associated tree. These rooted trees form a set $\mathcal{R}_n^*$. Notice that $|\mathcal{R}_n| \sim \frac{D \cdot x_0^{-n}}{n^{\frac{3}{2}}}$ and $|\mathcal{T}_n^*| \sim \frac{C \cdot x_0^{-n}}{n^{\frac{5}{2}}}$. We get $\frac{|\mathcal{R}_n^*|}{|\mathcal{R}_n|} \rightarrow 0$. In conjunction with Theorem 3, we have that the number of vertex classes is $(\mu_r + o(1))n$ for almost all rooted trees in $\mathcal{R}_n^*$.

According to whether the number of vertex classes is $(\mu_r + o(1))n$ or not, we depart $\mathcal{R}_n^*$ into two parts $\mathcal{R}_{n,1}^*$ and $\mathcal{R}_{n,2}^*$. There are at most $\frac{|\mathcal{R}_{n,2}^*|}{\frac{1}{24}n}$ trees in $\mathcal{T}_n^*$ corresponding to $\mathcal{R}_{n,2}^*$. Since $|\mathcal{R}_{n,2}^*| = o(|\mathcal{R}_n^*|) = o(|\mathcal{R}_n|)$, then $\frac{|\mathcal{R}_{n,2}^*|}{\frac{1}{24}n} = o(|\mathcal{T}_n|) = o(|\mathcal{T}_n^*|)$.

Therefore, almost all trees in $\mathcal{T}_n^*$ correspond to the rooted trees in $\mathcal{R}_{n,1}^*$. And recall that the root of the tree in $\mathcal{R}_{n,1}^*$ is a fixed vertex. That is, almost all trees in $\mathcal{T}_n^*$ also have $(\mu_r + o(1))n$ vertex classes. Consequently, almost every tree in $\mathcal{T}_n$ has $(\mu_r + o(1))n$ vertex classes. The proof is complete. $\square$

From Theorem 5, we have an intuitive grasp that the rooted tree space is just the tree space with a scale $(\mu_r + o(1))n$. Not rigorously to say, if we consider any special structure in trees, the case that this structure will appear $(\mu_r + o(1))n$ times in rooted trees is in a large probability, and the probabilities of appearances in tree space and rooted tree space seem to be the same. Moreover, by the asymptotical values of $|\mathcal{R}_n|$ and $|\mathcal{T}_n|$, we can get that $\mu_r \approx 0.8210$.

4 The distribution for any pattern in $\mathcal{T}_n$

In this section, we shall focus on the distribution of the occurrences $X_{n,M}$ of a pattern $\mathcal{M}$ on tree space $\mathcal{T}_n$. It is known that the distribution of the occurrences of a pattern in

$\mathcal{R}_n$ is asymptotically normal. We refer the readers to [6]. We show that the corresponding distribution in $\mathcal{T}_n$ is also asymptotically normal. It has been shown that $X_{n,M}(T)$ has mean $(\mu_M + o(1))n$ and variance $(\sigma_M + o(1))n$ and for almost every tree, and the number of corresponding rooted trees is $(\mu_r + o(1))n$. The constants μ and σ are the same as those for the case of rooted trees, namely, $E(X_{n,M}(T)) \sim \mu_M n \sim E(X_{n,M}(R))$ and $Var(X_{n,M}(T)) \sim \sigma_M n \sim Var(X_{n,M}(R))$. Based on these two results, we proceed to get our final result.

Theorem 6. *For any given pattern, the number of occurrences of the pattern in trees is asymptotically normally distributed.*

Proof. Recall that for each given pattern $\mathcal{M}$, $E(X_{n,M}(T)) \sim \mu_M n$ and $Var(X_{n,M}(T)) \sim \sigma_M n$, where μ_M and σ_M are some constants. Let $\mathcal{T}_n^1$ be the subset of $\mathcal{T}_n$ such that the number of occurrences $X_{n,M}(T)$ satisfies that $\frac{X_{n,M}(T) - \mu_M n}{\sqrt{\sigma_M n}} \leq t$, where t is some real number. Then, the probability

$$P\left(\frac{X_{n,M}(T) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right) = \frac{|\mathcal{T}_n^1|}{|\mathcal{T}_n|},$$

where $\mathcal{T}_n^1$ is the subset of $\mathcal{T}_n$. For $\mathcal{R}_n$, we shall try to show that

$$\lim_{n \rightarrow \infty} P\left(\frac{X_{n,M}(R) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right) = \lim_{n \rightarrow \infty} P\left(\frac{X_{n,M}(T) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right).$$

We knew that

$$\lim_{n \rightarrow \infty} P\left(\frac{X_{n,M}(R) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right) = N(0, 1, t),$$

where $N(0, 1, t)$ denotes the probability value of the normal distribution at t . Denote by $\mathcal{R}_n^1$ the set of rooted trees satisfying $\frac{X_{n,M}(R) - \mu_M n}{\sqrt{\sigma_M n}} \leq t$. The last equation holds from the fact that any pattern in $\mathcal{R}_n$ is asymptotically normally distributed.

If R is a rooted tree in $\mathcal{R}_n^1$ corresponding to $T \in \mathcal{T}_n$, then $X_{n,M}(R) = X_{n,M}(T)$. So, a tree T is in $\mathcal{T}_n^1$ if and only if all the associated rooted trees are in $\mathcal{R}_n^1$. We depart $\mathcal{T}_n^1$ into two subsets, $\mathcal{T}_n'^1$ and $\mathcal{T}_n''^1$, one is the collection of trees corresponding to $(\mu_r + o(1))n$ rooted trees, and the other is not, respectively. By Theorem 5, the number of rooted trees corresponding to $\mathcal{T}_n'^1$ is $|\mathcal{T}_n'^1| \cdot (\mu_r + o(1))n$, and $|\mathcal{T}_n''^1| = o(|\mathcal{T}_n^1|)$, i.e., the number of rooted trees associated with $\mathcal{T}_n''^1$ is at most $o(|\mathcal{T}_n^1|) \cdot n$. Then, it follows that

$$|\mathcal{T}_n'^1| \cdot (\mu_r + o(1))n \leq |\mathcal{R}_n^1| \leq |\mathcal{T}_n'^1| \cdot (\mu_r + o(1))n + o(|\mathcal{T}_n^1|) \cdot n.$$

Since $o(|\mathcal{T}_n^1|) \cdot n = o(|\mathcal{R}_n^1|)$ and $\frac{|\mathcal{T}_n'^1|}{|\mathcal{T}_n^1|} \sim 1$, we have $|\mathcal{R}_n^1| = (\mu_r + o(1))n \cdot |\mathcal{T}_n^1|$.

Therefore, we get that

$$\begin{aligned} P\left(\frac{X_{n,M}(R) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right) &= \frac{\mathcal{R}_n^1}{\mathcal{R}_n} \\ &\sim \frac{\mathcal{T}_n^1}{\mathcal{T}_n} \\ &= P\left(\frac{X_{n,M}(T) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right). \end{aligned}$$

Consequently,

$$\begin{aligned} \lim_{n \rightarrow \infty} P\left(\frac{X_{n,M}(T) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right) &= \lim_{n \rightarrow \infty} P\left(\frac{X_{n,M}(R) - \mu_M n}{\sqrt{\sigma_M n}} \leq t\right) \\ &= N(0, 1, t). \end{aligned}$$

Then the variable $X_{n,M}(T)$ is also asymptotically normal with mean $\sim \mu_M n$ and variance $\sim \sigma_M n$. $\square$

Now, we have established that for any pattern, the limiting distribution of the number of occurrences in $\mathcal{T}_n$ is also normal, which solves an open question claimed in [7].

References

- [1] B. Bollobás, Random Graphs (2nd Ed.), Cambridge Studies in Advanced Math., Vol.73, Cambridge University Press, Cambridge, 2001.
- [2] F. Chyzak, M. Drmota, T. Klausner, G. Kok, The distribution of patterns in random trees, *Comb. Probab. Comp.* **17**(2008), 21–59.
- [3] M. Drmota, Systems of functional equations, *Random Structures Algorithms* **10**(1997), 103–124.
- [4] M. Drmota, B. Gittenberger, The distribution of nodes of given degree in random trees, *J. Graph Theory* **31**(1999), 227–253.
- [5] F. Harary, E.M. Palmer, Graphical Enumeration, Academic Press, New York and London, 1973.
- [6] G. Kok, Pattern distribution in various types of random trees, In: Proc. 2005 International Conference on Analysis of Algorithms, 223–230.

- [7] G. Kok, The distribution of patterns in random trees. Thesis, Institut für Diskrete Mathematik und Geometrie, TU Wien, Austria, 2005.
- [8] X. Li, Y. Li, The asymptotic value of the Randic index for trees, *Adv. Appl. Math.* **47**(2011), 365–378.
- [9] R. Otter, The number of trees, *Ann. Math* **49**(1948), 583–599.
- [10] G. Pólya, Kombinatorische Anzahbestimmungen für Cruppen, Graphen und chemische Verbindungen, *Acta Math* **68**(1937), 145–254.
- [11] R.W. Robison, A.J. Schwenk, The distribution of degrees in a large random tree, *Discrete Math.* **12**(1975), 359–372.